



COPING WITH QUITTING: COGNITIVE AND BEHAVIORAL STRATEGIES

| | |
|---|---|
| <p>COGNITIVE STRATEGIES focus on retraining the way a patient thinks. Often, patients will deliberate on the fact that they are thinking about a cigarette, and this leads to relapse. Patients must recognize that thinking about a cigarette doesn't mean they need to have one.</p> | |
| REVIEW COMMITMENT TO QUIT | Each morning, say, "I am proud that I made it through another day without tobacco!" Remind oneself that cravings and temptations are temporary and will pass. Announce, either silently or aloud, "I am a nonsmoker, and the temptation will pass." |
| DISTRACTIVE THINKING | Use deliberate, immediate refocusing of thinking toward other thoughts when cued by thoughts about tobacco use. |
| POSITIVE SELF-TALKS, PEP TALKS | Say, "I can do this," and remind oneself of previous difficult situations in which tobacco use was avoided. |
| RELAXATION THROUGH IMAGERY | Center mind toward positive, relaxing thoughts. |
| MENTAL REHEARSAL, VISUALIZATION | Prepare for situations that might arise by envisioning how best to handle them. For example, envision what would happen if offered a cigarette by a friend—mentally craft and rehearse a response, and perhaps even practice it by saying it aloud. |
| <p>BEHAVIORAL STRATEGIES involve specific actions to reduce risk for relapse. These strategies should be considered prior to quitting, after determining patient-specific triggers and routines or situations associated with tobacco use. Below are strategies for several of the more common cues or causes for relapse.</p> | |
| STRESS | Anticipate upcoming challenges at work, at school, or in personal life. Develop a substitute plan for tobacco use during times of stress (e.g., use deep breathing, take a break or leave the situation, call a supportive friend or family member, use nicotine replacement therapy). |
| ALCOHOL | <i>Drinking alcohol can lead to relapse.</i> Consider limiting or abstaining from alcohol during the early stages of quitting. |
| OTHER TOBACCO USERS | <i>Quitting is more difficult if the patient is around other tobacco users. This is especially difficult if another tobacco user is in the household.</i> During the early stages of quitting, limit prolonged contact with individuals who are using tobacco. Ask co-workers, friends, and housemates not to smoke or use tobacco in your presence. |
| ORAL GRATIFICATION NEEDS | Have nontobacco oral substitutes (e.g., gum, sugarless candy, straws, toothpicks, lip balm, toothbrush, nicotine replacement therapy, bottled water) readily available. |
| AUTOMATIC SMOKING ROUTINES | Anticipate routines associated with tobacco use and develop an alternative plan. Examples: MORNING COFFEE: change morning routine, take shower before drinking coffee, drink tea instead of coffee, take a brisk walk shortly after awakening. WHILE DRIVING: remove all tobacco from car, have car interior detailed, listen to an audio book or talk radio, use oral substitutes. WHILE ON THE PHONE: stand while talking, limit call duration, change phone location, keep hands occupied by doodling or sketching. WHILE WATCHING TV: sit in a different chair, rearrange furniture, consider watching in a different room, keep hands busy by squeezing a stress ball. AFTER MEALS: get up and immediately do dishes or take a brisk walk after eating, brush teeth, call supportive friend. |
| POST-CESSATION WEIGHT GAIN | Do not attempt to modify multiple behaviors at one time. If weight gain is a barrier to quitting, engage in regular physical activity and adhere to a healthful diet (as opposed to strict dieting). Carefully plan and prepare meals, increase fruit and water intake to create a feeling of fullness, and chew sugarless gum or eat sugarless candies. Consider use of pharmacotherapy shown to delay weight gain. |
| CRAVINGS FOR TOBACCO | Cravings for tobacco are temporary and usually pass within 5–10 minutes. Handle cravings through distractive thinking, take a break, do something else, take deep breaths. |